
1st Re–organization for the Step–by–step Daily Recovery (11.1~) Follow–up Response (11.29~)

- □ The 2nd re-organization is postponed in consideration of the pandemic trend, such as the rapid increase in the number of confirmed cases.
 - Promote a follow-up response plan for medical and quarantine while maintaining the current level for 4 weeks before the effect of booster vaccination.

□ Home-based treatment Transformation

 A system in which patients are admitted to a medical institution only in exceptional case where home-treatment is not possible.



• 24 hours consultation in case of emergency, establish a hotline for treatment, advance designation of a transfer medical institution

□ Encouraging non-vaccinated people and promptly implementing booster shots.

• Encouraging non-vaccinated people

- Resumption of advance reservations for children and adolescents(12~17 years of age) (~'22.01.22.), and vaccination through School-Vaccination Institution linkage depending on local conditions (organized by the Office of Education).
- Cooperating with the medical community and encouraging active vaccination of the elderly who often visit medical institutions for other treatment

• Implementation of Rapid Booster Vaccination

- In December is the operation for intensive vaccination period for Seniors (60 years old and up), similar to influenza vaccination, same-day vaccination is administered without prior reservation.
- For all those who have completed basic vaccination over the age of 18, booster vaccination is administered 5 months (150 days) after the basic vaccination
- Set the validity period (6 months) for the quarantine pass(vaccination certificate), and additional vaccination are required within 6 months to use facilities that require a quarantine pass.
- Individual condition (disease treatment, overseas departure, etc), group vaccinations (facilities that are vulnerable to infection, medical institutions, etc) and those wishing to receive residual vaccines can receive early vaccinations within 1 month compared to the standard^{*}standby for additional vaccinations by target group.

 * (4 months after the completion of basic vaccination) 60 years old and up, those with underlying disease, medical institutions, facilities vulnerable to infection

(5 months after the completion of basic vaccination) 18-49 years old, age of 50's

(2 months after the completion of basic vaccination) for those who are vaccinated with Janssen vaccine, for those who are of with weak immune system.

\Box Strengthen Quarantine

- Reinforcement of PCR testing* for facilities that are vulnerable to infection, such as Nursing Hospitals and facilities, and it is focused to those who have completed the additional vaccine (booster shot).
 - * (Standard) 1 PCR test (when hospitalized) + 1 day waiting in isolation room \rightarrow (Changes) 2 PCR tests (^①when hospitalized, in isolation room for 3rd day) + 4 days waiting in isolation room
 - Winter Kimchi events are prohibited at leisure welfare facilities for Elderly(Senior Center, Senior Welfare Center, etc.)

<This translation is made by Danuri Helpline Center1577-1366>